

X. ALTERNATIVES FOR DISPOSAL OF EXCAVATED MATERIAL

The restoration alternatives will require excavation volumes ranging from 3,600 to 120,000 cubic yards. A preliminary investigation of potential disposal locations was conducted to help evaluate the feasibility of this excavation. Figure X-1 shows the locations of the alternative disposal sites; these are summarized in Table X-1 and below in terms of capacity, and compatibility with existing land use. Final selection of a site will require further environmental analysis of the impacts of disposal on jurisdictional wetlands, riparian habitat, floodplain restoration, and upland habitats.

A. DISPOSAL BEYOND THE REDWOOD CREEK/GREEN GULCH WATERSHED

Disposal outside of the Redwood Creek/Green Gulch watershed would require long truck hauls over narrow roads such as Highway 1. In addition to the high disposal cost, this would involve extensive environmental impacts from the increased truck traffic. This alternative was therefore deemed infeasible for this preliminary assessment.

B. GREEN GULCH FIELDS 6 AND 7

These are the two lowest fields operated by the Green Gulch Farm, and would be adjacent to the restored wetlands. Field 6 is currently used intermittently for vegetable farming, and Field 7 is used for horse pasturing. Part of Field 7 would be included in the restored wetland.

The Green Gulch Farm has expressed interest in using fill material to level these fields and improve drainage (Rudnick, personal communication). Together Fields 6 and 7 would have about 4 acres available for disposal, with elevations ranging from +12 feet NGVD near the proposed wetland to +25 feet at the top of field 6. Excavated material could be placed to level most of the fields at between +20 and +26 feet NGVD (providing about a 1% slope for drainage). The fields could then slope down towards the edge of the restoration project at about 10:1. The fields could accommodate about 24,000 cubic yards of excavated material with this disposal configuration.

Important considerations for disposal include:

- The material should be tested for suitability as agricultural soil. Soil corings indicate that most of the material would consist of 1-2 feet of productive top soil underlain by several feet of silty sands. The silty sands would probably not be suitable as top soil, but could provide good drainage as substrate below either dredged top soil or stockpiled soil from the fields.

- Material handling to segregate and stockpile soil could increase costs of excavation and disposal.
- Disposal should provide adequate setback from Green Gulch Creek to accommodate future restoration efforts.
- Disposal should not interfere substantially with Green Gulch Farm's farming activities. The fields are currently not frequently used, and could be left out of production during excavation.
- Access would be directly through the fields and along an existing dirt road. Green Gulch is also interested in using suitable fill material to raise and stabilize this road. The longest truck haul would be about 1300 feet.

The 1.4 acre field below the upper reservoir was also considered but rejected because of adverse impacts on Green Gulch's leach field. Other sites in upper Green Gulch were not considered feasible, since these would require access through the main entrance and would unduly disrupt the functions of the Green Gulch Farm.

C. BANDUCCI FLOWER BULB AND HEATHER FARM

Banducci currently grows flowers on about 28 acres of level fields in Franks Valley adjacent to Redwood Creek. The fields are active for most of the year, except for the late fall and early winter months. Preliminary discussions with Banducci indicate that it is feasible to raise these fields by about 2 to 3 feet. Allowing for a 50 to 100-foot setback from Redwood Creek, this would leave 25 acres available for disposal. The fields could therefore accommodate between 80,000 and 120,000 cubic yards of fill. Important considerations for disposal include:

- The material should be tested for suitability as agricultural soil. Soil corings indicate that most of the material would consist of 1-2 feet of productive top soil underlain by several feet of silty sands. The silty sands would probably not be suitable as top soil, but could provide good drainage as substrate below either dredged top soil or stockpiled soil from Banducci's farm.
- Historically these fields probably functioned as the Redwood Creek floodplain during extreme floods. These fields have not been observed to flood since levees were constructed at selected low points (Banducci, 1993). Disposal on these fields should therefore account for potential future restoration of the historic floodplain.

One strategy would be to place the material at larger depths over a smaller area, and provide a wider setback from the creek bank.

- Excavation would probably occur in the summer, when the fields would be in full production. Banducci may therefore have to be compensated for loss of income during construction.
- Access would be on a dirt road that enters Highway 1 about 1800 feet from Pacific Way. This road would probably have to be raised and improved. There is potential to use excavated gravel and coarse material for this purpose. The longest truck haul would be about 5800 feet.

D. RIDING RING AND STATE PARK LAND ABOVE BANDUCCI

A riding ring and former dairy pasture cover about 7.5 acres in Franks Valley above Banducci's farm. Existing land uses include horse riding and hiking. This gently sloping area could probably be raised by 2 to 3 feet, providing 24,000 to 36,000 cubic yards of disposal. Important considerations include:

- Grading of the disposed material should be done to provide a natural topography that would not interfere with existing uses of the site.
- The material should be revegetated to restore existing meadow habitat values.
- The lower portions of the area are part of the Redwood Creek floodplain; adequate setbacks from the creek should be provided to minimize flood hazards and impacts on the geomorphic stability of the creek.
- Access would be through a fire road/trail that enters Franks Valley Road. The longest haul would be about 7500 feet.
- Much of the land is owned by the Mt. Tamalpais State Park. Disposal should be done in a manner consistent with the State Park's goals for the site.

E. THE BALLFIELD AREA IN LOWER FRANKS VALLEY

This area covers about 5.5 acres of fallow fields near the intersection of Franks Valley Road and Highway 1. The site is bordered by Franks Valley Road on south side and Redwood Creek on the northern side. A 2 to 5-foot levee separates the entire field from the creek; construction of this levee has been observed to cause increased flooding upstream (Banducci, personal communication).

Although this field is commonly referred to as the "Ballfield", it is currently overgrown with grasses and coyote brush. Raising the field by about 2 to 3 feet would provide 18,000 to 27,000 cubic yards of disposal. Important considerations include:

- Historically this field probably functioned as the Redwood Creek floodplain during extreme floods. These fields have not been observed to flood since the levees were constructed. Disposal should therefore account for potential future restoration of the historic floodplain.
- The site is owned by the GGNRA, and should be graded and revegetated in a manner consistent with Park Service goals for the site.
- Access would be directly off of Franks Valley Road, with a truck haul of about 7500 feet.

F. DISPOSAL ALONG THE PERIMETER OF THE RESTORED WETLAND

Upland areas adjacent to the restored wetland could be raised up to the level of the Highway 1. These would include the horse ring and Green Gulch Farm lots at the corner of Pacific Way and Highway 1, as well as the narrow area currently occupied by the Green Gulch trail along Highway 1.

The Highway currently lies at elevations between +17 and +19 feet NGVD. Assuming a 4:1 slope up from the edge of the restored riparian fringe, about 20,000 cubic yards could be disposed of on these uplands. Key considerations include:

- This would require minimal hauling, and would be the least expensive disposal alternative.
- The slope from the wetland up to the maximum fill elevation should be graded in a manner that provides a natural transition from the wetland to the road.

- The fill should be revegetated in a manner compatible with the restored wetland habitat, using non-invasive native plants.

XI. REFERENCES

- American Ornithologists' Union. 1983. Check-list of North American Birds. Sixth Edition. A.O.U., Washington, D.C.
- American Ornithologists' Union. 1985. Thirty-fifth supplement to Amer. Ornith. Union Check-list of North American Birds. *Auk* 102:680-686.
- Atwater, B. S. Conrad, T. Dowden, D. Hedel, R. MacDonald and W. Savage. 1979. History, landforms and vegetation of the estuary's tidal marshes. pp. 347-386. *In*: T. Conomos, editor, San Francisco Bay: the urbanized estuary. Pacific Division, AAA, San Francisco, CA.
- Banducci, Amadeo. 1993. Personal Communication. Manager of Flower Bulb and Heather Farm in Franks Valley.
- Barbour and Major. 1988. Terrestrial vegetation of California. Calif. Native Plant Soc. Sacramento, CA. Spec. Publ. No. 9. ix + 1030 p.
- Barbour, M., B. Pavlik, F. Drysdale and S. Lindstrom. 1993. California's changing landscapes diversity and conservation of California vegetation. Calif. Native Plant Soc. Sacramento, CA. 246 p.
- Barbour, M., R. Craig, F. Drysdale and M. Ghislen. 1973. Coastal ecology Bodega Head. Univ. Calif. Press, Berkeley, CA. xix + 338 p.
- Borror, D. J. and R. E. White. 1970. A Field Guide to Insect . Houghton Mifflin, Boston.
- California Department of Fish and Game. 1991. Bird and mammal Species of Special Concern. Sacramento.
- Carr, A. 1952. Handbook of Turtles. Comstock Publishing Asso. Cornell University Press. Ithaca, New York. 542p.
- Cowen, R. 1991. Vanishing amphibians: Why they're croaking. *Science News*. 137:116.
- Fitch, H.S. 1938. *Rana boyleii* in Oregon. *Copeia* 1938:148.

Duncan, Faith. date unknown. *The History of Two Valleys: Frank's and Homestead Valley, Marin County, California*. Muir Woods Library.

Evans, J.G. 1988. The natural history of the Point Reyes peninsula. Point Reyes Nat. Seashore Assoc., Point Reyes.

Federal Emergency Management Agency. Flood Insurance Rate Maps for Marin County.

Garcia, A.W., *et al.* 1975. Type 16 Flood Insurance Study: Tsunami Predictions for Monterey and San Francisco Bays and Puget Sound.

Goerke, Betty. 1993. Personal Communication. College of Marin Anthropology Department.

Gordon, B. 1987. Monterey Bay area: natural history and cultural imprints. 2nd Ed. Boxwood Press, Pacific Grove, CA.

Grinnell, J. and A.H. Miller. 1944. Distribution of the birds of California. *Pac. Coast Avifauna* 27.

Grinnell, J., H.C. Bryant and T.I. Storer. 1918. The game birds of California. Univ. Calif. Press., Berkeley.

Harding-Lawson and Associates. 1991. Water Quality Sampling In Redwood Creek. Letter Report prepared for the Muir Beach Community Services District.

Hayes, M. P. and M. R. Jennings. 1986. Decline of ranid frog species in western North America: Are bullfrogs (*Rana catesbeiana*) responsible? *J. Herp.* 20:490-509.

Hayes, M. P. and M. R. Jennings. 1989. Habitat correlates of distribution of the California red-legged frog (*Rana aurora draytonii*) and the foothill yellow-legged frog (*Rana boyleii*): Implication for management. In: Robert E. Sarzo, Keith E. Severson, and David R. Patton (technical coordinators): Management of amphibians, reptiles, and small mammals in North America, Proceedings of the symposium on the (July 19-21, 1988, Flagstaff, AZ). USDA General Technical Report RM-166:1-458.

Heard, M. 1904. A California frog ranch. *Out West* 21: 20-27.

Herbold, B. and P.B. Moyle. 1989. The ecology of the Sacramento-San Joaquin Delta: A community profile. *US Fish Wildl. Serv. Biol. Rep.* 85(7.22).

- Hickman, J. (ed.). 1993. The Jepson manual, higher plants of California. Univ. Calif. Press, Berkeley, CA. xvii + 1400 p.
- Hofstra, T.D. and D.G. Anderson. 1989. Survey of Salmonid Fish and Their Habitat. Redwood Creek, Marin County, California. Redwood National Park Technical Services Division, Arcata, California.
- Holland, R. 1986. Preliminary descriptions of the terrestrial natural communities of California. Non-game heritage program. Calif. Dept. Fish Game, Sacramento, CA. unpub. m.s. 156 p.
- Holland, R. 1990. Natural communities. Natural diversity data base. Calif. Dept. Fish Game, Sacramento, CA. unpub. m.s. 9 p.
- Hornberger, M. 1990. Paleoenvironment of Elkhorn Slough and surrounding wetland habitats: a geological study using and ecological approach. M.S. Thesis, Moss Landing Marine Laboratories, Moss Landing, CA.
- Howell, J. 1969. Marin flora, manual of flowering plants and ferns of Marin County, California. Univ. California Press, Berkeley, CA. viii + 366 p.
- Ingles, L. 1965. Mammals of the Pacific states. Stanford Univ. Press, Stanford, CA. xii + 506 p.
- Jameson, E. and H. Peeters. 1988. California mammals. California Natural History Guides, 52. Univ. California Press, Berkeley, CA. xi + 403 p.
- Jennings, M.R. 1991. Status of the California red-legged frog (*Rana aurora draytonii*) in Moore Creek on the University of California at Santa Cruz campus. unpubl. tech. rep.
- Jennings, M.R. and M.P. Hayes. 1985. Pre-1900 overharvest of California red-legged frogs (*Rana aurora draytonii*): The inducement for bullfrog (*Rana catesbeiana*) introduction.
- Johns, E.L. 1989. Water Use by Naturally-Occurring Vegetation, Including an Annotated Bibliography. ASCE Irrigation and Drainage Division.
- Johnson, J.W. 1973. Characteristics and Behavior of Pacific Coast Tidal Inlets. Journal of Waterways, Harbors, and Coastal Engineering Division. ASCE. Vol. 99, No. WWE. August. pp. 325-339.

Lehre, A.K. 1974. The Climate and Hydrology of the Golden Gate National Recreation Area. From "The Terrestrial Environment of the Golden Gate National Recreation Area". Prepared by the University of California at Berkeley for the National Park Service.

Lehmkuhl, D. M. 1979. Aquatic Insects. Wm. C. Brown Company, Dubuque.

Mansell, W. 1980. North American birds of prey. Morrow, Williams & Co. Inc., New York.

Moyle, P.B. 1973. Effects of introduced bullfrogs, *Rana catesbeiana*, on the native frogs of the San Joaquin Valley, California. *Copeia*. 1: 18-22.

Munro-Fraser, J.P. 1880. *The History of Marin County, California*. Alley, Bowen and Company Publishers, San Francisco.

Myers, Robin. 1990. Redwood Creek Water Balance, Hydrologic Year 1985-86. Muir Woods National Monument. November.

Natioanl Geographic Society. 1983. Field Guide to the birds of North America. National Geographic Society, Washington, D.C.

National Research Council, Marine Board. 1987. Responding to Changes in Sea Level: Engineering Implications. National Academy Press, Washington, D.C.

Omduff, R. 1974. Introduction to California plant life. University California Press, Berkeley, CA. 152 p.

Pechmann, J.H.K., D.E. Scott, R. D. Semlitsch, J.P. Caldwell, L.J. Vitt, and J. W. Gibbons. 1991. Declining amphibian populations: The problem of separating human impacts from natural fluctuations. *Science*. 253: 892-895.

Philip Williams & Associates, Ltd. 1994. Summary of Findings on the Impacts of the Banducci Flower and Bulb Farm. Technical Memorandum to Nancy Horner, Golden Gate National Recreation Area, San Francisco, CA.

Phillips, K. 1990. Where have all the frogs and toads gone? *Bioscience*. 40: 1-6.

Pickwell, G. 1949. Amphibians and Reptiles of the Pacific States. Stanford University Press. Stanford, California. 236p.

Powell, J. A. and C. L. Hogue. 1979. California Insects. University of California Press, Berkeley.
Rathbun, G., N. Siepel, and D. Holland. 1992. Nesting behavior and movements of western pond turtles, *Clemmys marmorata*. Sw. Nat. 37(3):319-324.

Roundtree, R. 1973. A morphological change in a California estuary: sedimentation and marsh invasion at Bolinas Lagoon. PH.D. Dissertation, University of California, Berkeley, CA.

Rudnick, Peter. 1993. Personal Communication. Green Gulch Farm.

San Francisco Bay Regional Water Quality Control Board. 1990. Update to the San Francisco Bay Basin Plan.

Schoenherr, A. 1992. A natural history of California. California Natural History Guides: 56. Univ. California Press, Berkeley, CA. xi + 772 p.

Schwartz, D., H. Mullins and D. Bellknap. 1986. Holocene geologic history of a transform margin estuary: Elkhorn Slough, California. Estuarine, Coastal and Shelf Science 22: 285-302.

Shuford, W. 1993. The Marin County breeding bird atlas: a distributional and natural history of coastal California birds. Bushtit Books, Bolinas.

Shuford, W. and I. Timossi. 1989. Plant communities of Marin County. Calif. Native Plant Soc. Sacramento, CA. Spec. Publ. No. 10, 32 p.

Shuford, W., G. Page, J. Evens, and L. Stenzel. 1989. Seasonal abundance of waterbirds at Point Reyes: a coastal California perspective. Western Birds 20(4): 137-265.

Skinner, J.E. 1962. A historical review of the fish and wildlife resource so the San Francisoc Bay ara. Calif. Dept. Fish and Game, Water. Proj. Branch Rep. 1.

Small, A. 1974. The birds of California.

Smith, J. and K. Berg. 1988. Inventory of rare and endangered vascular plants of California. Calif. Native Plant Soc. Sacramento, CA. Spec. Publ. 1 (4th edition), 246 p.

Soil Conservation Service. 1979. Soil Survey of Marin County, California.

Sousa, Joseph. 1993. Personal Communication. Former Dairyman along Redwood Creek.

Stebbins, R.C. 1954. Amphibians and Reptiles of Western North America. McGraw-Hill Book Company, Inc. New York, New York. 536p.

Storm, R.M. 1960. Notes on the breeding biology of the red-legged frog (*Rana aurora aurora*) Herpetologica 16: 251-259.

U.S. Army Corps of Engineers. Wetland Delineation Manual. U.S. Department of Commerce.

U.S. Army Corps of Engineers. 1987. Corps of Engineers Wetland Delineation Manual, Department of the Army Technical Report Y-87-1.

U.S. Army Corps of Engineers. 1965. Cooperative Beach Erosion Study: Pt. Delgado to Pt. Ano Nuevo, California. San Francisco District.

U.S. Coast and Geodetic Survey. 1853. Pacific Coast, North of the Entrance to San Francisco Bay. Topographic Sheet 400, March.

U.S. Fish and Wildlife Service. 1987. Migratory nongame birds of management concern in the United States: The 1987 list. Office Migratory Bird Mgmt, USFWS, Washington, DC.

U.S. Geological Survey. 1894. Tamalpais Sheet.

U.S. Geological Survey. 1970. Surface Water Supply of the United States, 1961-1970. Water Supply Papers 1929 and 2129.

Zedler, J. 1984. Salt marsh restoration: a guidebook for southern California. California Sea Grant Report No. T-CSGCP-009.

Zeiner, D., W. Laudenslayer, K. Mayer and M. White. 1990. California's wildlife, Volume III, mammals. Calif. Dept. Fish and Game. Sacramento, CA. xi + 407 p.

Zweifel, R.G. 1955. Ecology, distribution, and systematics of frogs of the *Rana boylei* group. University of California Publications in Zoology 54(4):207-292.

ACKNOWLEDGEMENTS

We would like to thank the following individuals for their assistance in preparing this report:

Amadeo Banducci	
Gordon Bennett	
Tony Brazil	
Kent Dedrick	
Renee Dixon	Caltrans
Glenn Fuller	Muir Woods National Monument
Betty Goerke	College of Marin Anthropology Department
Daphne Hatch	Golden Gate National Recreation Area
Nancy Hornor	Golden Gate National Recreation Area
Henry Hyde	Muir Beach Community Services District
Eric Larson	Hydrologist
Mia Monroe	Muir Woods National Monument
Peter Rudnick	Green Gulch Farm
Yvonne Rand	Green Gulch Farm
Sid Shadle	Caltrans
Nancy Skinner	Naturalist
Joseph Sousa	
Terri Thomas	Golden Gate National Recreation Area
Ed Ueber	National Marine Sanctuary
Denise Vore	Golden Gate National Recreation Area
Dr. Lisa Wells	University of California, Berkeley
George Wheelwright	

The primary authors of this report were:

Robert Schanz	Project Manager, Philip Williams & Associates, Ltd.
Philip Williams	Principal-in-Charge, Philip Williams & Associates, Ltd.
Jim Oakden	Project Manager, Moss Landing Marine Labs
Dr. Jerry Smith	San Jose State University
John Roberts	John Northmore Roberts and Associates
Nancy Hornor	Golden Gate National Recreation Area

The following individuals also contributed to the report:

Andrea Bennett	Philip Williams & Associates, Ltd.
Tamara Rose	Philip Williams & Associates, Ltd.
John Oliver	Moss Landing Marine Laboratories
Peter Slattery	Moss Landing Marine Laboratories
Diane Carney	Moss Landing Marine Laboratories
Jo Guerrero	Moss Landing Marine Laboratories
Ed Ely	Aquatic and Wildlife Biologist
